

**Effects of Cover Crop and Nitrogen Rate on Corn Grain and Silage Yield,
Nitrogen Loss in Tile Drainage and Soil Health.
PROJECT NUMBER: R2023-13 MN DEPT OF AGRICULTURE
Award: CON000000102844
2023 Annual Report**

Jeffrey Vetsch
Univ. of Minnesota
Southern Research and Outreach Center
Waseca, MN 56093-4521

Introduction / Justification

Nitrogen is an essential input for profitable corn production. Previous research has shown subsurface tile drainage systems deliver nitrate-N to surface waters and thereby degrade water quality (Randall and Mulla, 2001, Dinnes et al., 2002). Row crop agriculture in the Midwest is under scrutiny to reduce nitrate concentrations and loads in tile drainage. The use of cover crops and applying appropriate rates of N for corn are potential management strategies to reduce nitrate loads in tile drainage water. However, research in Minnesota has shown cover crop establishment can be difficult (Strock et al., 2004), often producing minimal cover crop growth (biomass) which results in less or inconsistent nitrate reduction in tile drainage water compared to other areas in the Midwest with longer growing seasons and milder winters (Kaspar et al., 2007).

There are cropping systems where cover crops could be more effective as nitrate scavengers and soil protectors. These crop systems include following canning crops, like sweet corn and peas, small grains and when corn is harvested for silage in early September. In these systems there is considerably more time for cover crop establishment and growth in the fall before soils freeze in Minnesota. Furthermore, after silage corn a cover crop could protect the soil from erosion and potentially replenish carbon lost during the silage (biomass) harvest which would improve sustainability.

The species of cover crop, establishment date and termination date can affect its potential to sequester N and carbon (C). Cereal rye is effective at scavenging N when it's established early and not terminated until spring. However, Vetsch et al. (unpublished) found cereal rye can negatively affect corn production and economics by decreasing yield or increasing input costs due to greater fertilizer N requirement, cover crop seed costs and herbicide costs for cover crop termination. A cover crop blend like oat, forage pea and radish are less expensive alternatives than cereal rye due to seed costs and no herbicide needed for termination. The potential of winter terminated cover crops to scavenge N and sequester C in a corn silage crop system in Minnesota is not known. Furthermore, it's not clear if nitrate loss in tile drainage is different between corn silage and corn grain systems.

Farmers are interested in the soil health benefits of cover crops, but also their potential to sequester carbon, especially in continuous corn systems with minimal or reduced tillage. The proposed study, which was initiated

in 2021 on the tile drainage research facility in Waseca, provides an opportunity to simultaneously measure the effects of cover crops on corn production, nitrate loss in tile drainage and soil health metrics.

The objectives of this study are to quantify the effects and interactions of cover crops and N rates on corn production, nitrate-N concentration and loss in tile drainage, N uptake, NUE, economic return and soil health parameters.

Experimental Procedures

A research experiment was initiated in 2021 at the Univ. of Minnesota Southern Research and Outreach Center drainage research facility on a poorly drained Canisteo-Webster clay loam soil complex. Thirty-six individual tile drainage plots were installed in 1976. Each plot, measures 20 ft. by 30 ft., has a separate drain outlet and is isolated from adjacent plots to minimize lateral flow. A single tile is placed four ft deep perpendicular to the rows. The plot spacing simulates a 50-ft. tile drain spacing. A randomized complete block design with 4 replications was used in this study. A restriction on randomization within blocks, based on previous tile flow history, helped balance variability in tile flow among the 36 plots. This restriction puts plots with the greatest historical flow all in the same block.

A total of 11 treatments were included in this study. Eight treatments were comprised from a partial factorial combination of three management factors: corn crop system (corn for grain and corn silage), cover crop use and N rate. The three cover crop treatments included: no cover crop in both crop systems (corn for grain and corn silage), cereal rye with spring termination or a blend of annuals (oat, forage pea and radish) with late fall or winter termination due to freezing. The four crop system treatments were corn for grain no cover crop (Gnc), corn for silage no cover crop (Snc), corn for silage with cereal rye cover (Srye) and corn for silage with annual blend cover (Sblend). These cover crop treatments were no-till drilled soon after corn silage harvest and only in the corn silage crop system. Cover crop seeding rates were 60 lb/ac for cereal rye and 18, 8, and 1 lb/ac for oat, forage pea, and radish, respectively. Nitrogen rates of 180 and 220 lb N/ac for continuous corn were compared across all crop systems and cover crop treatments. The 180-lb rate is near the maximum of the MRTN Profitable N Rate Range of 159 to 189 lb/ac for corn after corn in Minnesota (0.10 price ratio) from the [Corn N Rate Calculator](#). The 220-lb rate aids in determining a cover crops ability to sequester N thereby minimizing potential nitrate loss in tile drainage water. Three additional N rate treatments were included in the study. One was a “zero” N control, which received 4.6 lb N/ac from liquid starter fertilizer, and the others were 140 and 260 lb N/ac for corn grain production. These additional treatments help in determining the optimum N rate for corn grain production. The 140 and 260 lb N/ac treatments did not have tile drainage monitoring.

Nitrogen fertilizer treatments were split-applied with 20 lb N/ac at planting and the remainder applied at V2 as urea ammonium nitrate (UAN, 32-0-0) which was stream-injected about 10 inches from the row. Liquid starter fertilizer, ammonium poly phosphate (APP, 10-34-0 at 4 gal/ac; 4.6 lb N/ac + 16 lb/ac of P₂O₅), was applied in-furrow at planting to all plots. All treatments, except for the control, received 3 gal/ac of UAN (10.7 lb N/ac) and

3.5 gal/ac of ammonium thiosulfate (ATS, 12-0-0-26S; 4.7 lb N/ac + 10 lb S/ac) in a surface-dribbled band three inches from the corn row at planting. See table 1 for completion dates of each procedure and for each year. See Supplemental Materials Table S1 for products used, rates applied and other information.

Corn was planted at 34,000 seeds/ac in 30-inch rows. The cereal rye treatments received a broadcast-application of glyphosate to terminate the cover crop, a second application of glyphosate was required in 2022 due to unusually cold weather which reduced the effectiveness of the first application. Additional weed control included pre and post emergence applications at full labeled rates (Table S1). Corn plant counts were taken from harvest rows. Relative leaf chlorophyll content (RLC) was calculated from Minolta SPAD meter measurements from the ear leaf (30 measurements per plot) at R1. Plot notes (corn height and greenness differences) were taken at the V8 and R3 corn growth stages. Corn silage yields were determined by hand harvesting two corn rows 10 ft in length. Plants were cut 8 inches above the soil surface (same as commercial harvesting). Whole plot samples were weighed wet, then four plants were chopped for a subsample to determine harvest moisture and this sample was dried at 140° F for 3 days, ground and submitted to a commercial lab to determine feed value (NDF, ADF, etc.) and nutrient content in corn silage. The remaining silage crop was removed with a custom harvester and cover crops were seeded with a no-till drill.

Corn grain yield from select treatments was harvested with a plot combine (two rows 33 ft in length). A grain sample was collected at harvest and this sample was dried at 140° F for 3 days, ground to and analyzed for nutrient content after microwave acid digestion at a commercial lab. Nitrogen removal in corn grain was calculated from grain yield and N concentration data. Nitrogen use efficiency parameters: partial factor productivity, PFP (the ratio of the grain yield to the applied rate of N) and agronomic efficiency, (the ratio of the increase in grain yield over N-control plots to the applied rate of N) were calculated as described by Snyder and Bruulsema (2007). For these NUE calculations the 4.6 lb N/ac rate from starter fertilizer was assumed to be the zero N control.

Cover crop biomass yields were measured by cutting and collecting all material from two 20 by 30 inches (4.17 sq. ft) areas from each plot in the fall and prior to termination in spring. No biomass harvest was conducted in the fall of 2022 due to very little cover crop growth (Appendix Pic. 5). A very dry September and October contributed to the poor cover crop growth in 2022. Since the annual blend cover terminated during the winter, these plots were not sampled in spring. Biomass samples were dried at 140° F for 3 days, weighed, ground, and analyzed for C and N concentration at a commercial lab. A corn stover (residue) sample was collected from the corn grain production plots, these samples were dried at 140° F for 3 days, weighed, ground, and analyzed for C and N concentration.

Corn stalks were chopped (shredded) about 6 inches above the soil surface on corn grain plots prior to fall strip tillage. All plots were strip tilled to an 8-inch depth with a Redball™ Model 2000. A fertilizer blend of P-K-S (Table S1) was applied in the strip-till band.

Tile drainage is measured via an automated collection system. Tile water collects in drainage wells, then is pumped via a sump pump through water meters that measure flow volume. Flow volume is recorded on a datalogger hourly. These hourly flow data are examined for outliers prior to summarizing daily. The previous 24-hours of flow are summed at 8 am each day. Whenever the sump pump turns on and pressurizes the system, a portion of the flow is collected in containers. This insures some of the water sample comes from the entire period between sample collection. This technique is often referred to flow-weighted sampling (Dinnes et al., 2002). Tile water samples are taken from each plot once a week during normal tile flow and two times per week during heavy tile flow. Water samples are kept cool prior to collection and then frozen after collection.

Each year soil samples were taken from all plots in the spring prior to planting corn at 0- to 15-, 15- to 30-, 30- to 60-, and 60- to 90-cm depths. Three 1.5-inch diameter (3.8 cm) cores are taken per plot. One core from a non-wheel track row is used to determine soil bulk density. The other cores are composited, immediately dried at 105° F, then ground and sieved to pass a 2-mm screen. Spring 0- to 15-cm samples were analyzed for Olsen P and exchangeable K using standard soil test methods for the North Central Region. All spring samples were analyzed for nitrate-N, ammonium-N, total N, total organic C, POXC and CO₂ burst (min-C) using standard soil test methods for the North Central Region. In November, fall soil samples were collected from a 0- to 30-cm depth, immediately dried at 105° F, then ground and sieved to pass a 2-mm screen. These fall samples were analyzed for nitrate and ammonium-N. The POXC and CO₂ burst tests were conducted in Dr. Anna Cates lab at the University of Minnesota. All other soil tests were conducted at commercial labs. These soil health tests allow us to assess microbial activity and food source. These C pools can be seen as early indicators of C sequestration and correlate with crop yields (Oldfield et al. 2021).

All data were statistically analyzed using ANOVA with Proc mixed in SAS® (SAS 9.4, SAS Institute Inc., 2014. Cary, North Carolina) after examination of residuals, outliers and normality assumptions using Proc univariate in SAS. A two-factor ANOVA with a split-plot arrangement of treatments compared the effects and interactions of crop system and cover crop treatments [grain corn, no cover crop; silage corn, no cover; silage corn, cereal rye with spring termination; and silage corn with annual blend with winter termination] and total N rate (180 and 220 lb/ac). Mean separations were determined using the P Diffs procedure in SAS with alpha=0.10 level of significance. Treatments followed by different letters within a row or column are significantly different.

Considerable variation in tile flow among individual plots is common in drainage research facilities. Managing it with blocking, statistical designs and replication is somewhat effective. Tile flow variability among treatments and among replications within a treatment often result in data that are not normally distributed and have non

constant variance among treatments. These assumptions are often violated and statistical analyses with ANOVA is not recommended. Furthermore, variability in tile flow among plots greatly influences nitrate loss or load among individual plots and treatment means. Thus, nitrate loss data are also not normally distributed and have non constant variance. Adjusting or correcting these data for flow during a period of collection (three month or annual periods) often results in data that are normally distributed and have constant variance. These techniques were used in the analysis of the data in this study and will be discussed further in the results and discussion section.

Results and Discussion

Weather

Weather data characterizing the 2021 growing season at Waseca are presented in Table 2. The data were taken from the SROC weather station (44.07064, -93.52645) located 0.3 miles from the drainage research site. The growing season was warm and dry as every month (April through September) had greater than normal mean temperatures and all months except August had considerably less than normal precipitation. An unusually dry period from April through mid-May (<0.9 inches of total precipitation), resulted in slow germination and uneven emergence of corn. The dry weather combined with liquid starter fertilizer injury of corn seedlings, a result of human error and equipment malfunction, required corn to be replanted on 24 May to minimize unevenness in corn stand and growth. Nitrogen loss due to leaching and denitrification would have been nearly zero during this droughty growing season as only two rainfall events of more than 1.0 inches occurred from April through September. Precipitation for the period from May through September totaled only 14.13 inches compared to the normal of 23.72 inches. Growing Degree Units (GDU, base 50 min and 86 max) totaled 2,979 a record for Waseca and 470 more than normal. Despite the dry conditions, corn yields were near long-term averages in 2021 which is remarkable.

Weather data in 2022 are presented in Table 2 and Figure 1a. Spring started cool and wet but turned warm and dry as May through September had greater than normal mean temperatures and June, July and September had less than normal precipitation. The cool and wet April and early May delayed corn planting and other field operations. Precipitation for the period from May through September totaled 19.98 inches compared to the normal of 23.72 inches. While monthly rainfall was near normal (± 0.7 inches) in all months except September, rainfall distribution was irregular. Each month had a large rainfall event including 2.41 inches on May 11-12, 3.50 inches on June 13-14 (2.5 inches in 2 hours), 2.50 inches on July 24, 1.93 inches on August 7-8 and 1.98 inches on August 28 (Figure 1b). The May and June events resulted in significant runoff and ponding in low areas of fields. Nitrogen loss from leaching was likely minimal due to rainfall intensity and runoff; however, leaching and denitrification in ponded areas likely occurred during these May and June events. These large rainfall events contributed about half of the total rainfall for a month. Furthermore, June, July, August and September had long dry periods with minimal or no rainfall. Growing Degree Units (GDU)

totaled 2,629, about 5% greater than normal. Despite this irregular weather, corn yields were only 10-15 bu less than average in 2022.

Weather data in 2023 are presented in Table 2 and Figure 1b. February, March and April were cool and wet which resulted in a heavy snowpack and flooded fields in March. Six inches of rainfall was recorded from May 6 through 15. Generally, the rest of May and June were very warm and very dry. May, June, August and September all had greater than normal mean temperatures while April and July were slightly cooler than normal. Summer months were quite dry as June through September precipitation totaled only 8.63 inches or 10.59 inches less than normal. The heavy mid-May rainfall event resulted in heavy tile flow and some culvert flooding. However, N loss from leaching and denitrification was minimal due to sidedress N application in early June. Growing Degree Units totaled 2,783 through September and 2,932 for the year or 17% greater than normal (2nd most all time). October and December had greater than normal precipitation with most of the December precipitation falling as rainfall on unfrozen soil which is very unusual in southern Minnesota. The cool and wet April and early May followed by a hot and very dry summer affected corn production. Corn grain and silage yields were 20% less than average in 2023.

Cover crop biomass yield

Treatments significantly affected nutrient concentration and uptake in cover crops and the C:N ratio (Table 3a). For the 5 November 2021 sampling, biomass yields were numerically greater with rye (296 lb/ac) than with blend (255 lb/ac) but not significantly greater (P value = 0.114), when averaged across N rates for corn. The blend had greater N concentration in biomass and a lower C:N ratio than rye. Carbon concentration, N uptake and C uptake were not affected by treatments and no significant interactions among treatment effects were observed. When averaged across cover crop species, the 170 lb N rate for corn in 2021 resulted in greater N concentration and uptake in biomass and lower C concentration and C:N ratio compared with the 140 lb N rate. These data suggest the 170 lb N/ac treatment had greater residual N, likely as NO_3^- , in the soil profile after harvest. However, the difference in N uptake in the cover crop biomass between the two N rates for corn was quite small, only 2 lb/ac.

Rye biomass yield averaged 296 lb/ac on 27 April 2022 compared with 276 lb/ac on 5 November 2021 (Table 3a). This small difference in biomass from fall to spring was not related to poor growth as rye height in April was about 2X greater than in November. The small increase in rye biomass from fall to spring was attributed to rye stand loss due to tractor wheel tracks and strip-tillage zones. Nitrogen rates for corn in 2021 had no effect on biomass yield and N and C concentration and uptake. However, C:N ratio was slightly less with 170 lb N/ac than with 140 lb N/ac.

Due to very poor growth during the dry fall of 2022 no cover crop biomass yields were taken in the fall of 2022. Rye biomass yields were collected on 8 May 2023 even though growth was minimal (Table 3b). Biomass

yields, nutrient concentrations and uptake were not affected by treatments. Rye biomass yield averaged less than 30 lb/ac of dry matter.

Treatments significantly affected cover crop biomass yield and nutrient uptake for the 23 October 2023 sampling (Table 3c). Cover crop biomass yield and nutrient uptake was much greater with blend (229 lb/ac) than with rye (92 lb/ac), when averaged across N rates for corn. A significant interaction among treatment effects was found for N concentration and C:N ratio. Nitrogen concentration was less with 220 lb N/ac than with 180 lb N/ac for rye but N concentration was greater with 220 lb N/ac than with 180 lb N/ac with blend. Moreover, the significant interaction for C:N ratio was a result of the interaction for N concentration, but the response was reversed due to N concentration being the denominator in the C:N ratio. The authors have no explanation for N concentration in the rye cover being less with a greater N rate for corn. We would expect it to be numerically greater with the higher N rate as it was with the blend. The 220 lb N/ac rate would likely have greater residual soil nitrate for the cover to scavenge.

Corn grain production

Corn grain yield and moisture from 2021 (study setup year) are presented in Table 4a. Grain yields ranged from 111 bu/ac in the control (4.6 lb N/ac) to 167 bu/ac with 170 lb N/ac for corn following soybean. An ANOVA analysis showed yields were statistically equal among the 140, 170 and 200 lb N/ac treatments. Grain moisture ranged from 20.0 to 21.2 percent and only small differences were observed as the control treatment was slightly drier.

Corn grain yield, moisture, nutrient concentration and nutrient uptake in 2022 are presented in Table 4b. Grain yields ranged from 60 bu/ac in the control (4.6 lb N/ac) to 218 bu/ac with 260 lb N/ac for corn following corn. ANOVA showed yields were greatest with 260 lb N/ac but statistically equal among the 140, 180 and 220 lb N/ac treatments. Grain moisture ranged from 22.7 to 26.2 percent. Moisture was least with 180 and 220 lb N/ac and greatest with 260 lb N/ac suggesting the higher N rate enhanced plant stay green, which may have contributed to slightly greater yields in this droughty growing season. Grain N concentration was least with the control and was statistically equal among other N rates for corn. Nitrogen removal in grain increased with increasing N rate and was statistically similar between the 220 and 260 lb N/ac rates. Grain P concentration was greatest in the control and numerically least with 260 lb N/ac. Grain K concentration was least with 260 lb N/ac and greatest with the control and the 180 and 220 lb/ac N rates. Grain S concentration was least with 260 lb N/ac and generally similar among other N rates. Grain P₂O₅ and K₂O removal was least with the zero N control and statistically equal among other N rate treatments. Grain removal averaged about 55 and 38 lb/ac of P₂O₅ and K₂O, respectively in the highest yielding treatments. Grain S removal was least with the zero N control and slightly less with 140 lb N/ac compared with other higher N rates.

Corn grain yield, moisture, nutrient concentration and nutrient uptake in 2023 are presented in Table 4c. Grain yields ranged from 48 bu/ac in the control (4.6 lb N/ac) to 163 bu/ac with 260 lb N/ac for corn following corn. The ANOVA showed yields were greatest with 260 lb N/ac but statistically equal to the 220 lb N/ac treatment. Grain moisture ranged from 19.4 to 21.7 percent. Moisture was least with 180 and 220 lb N/ac and greatest with the control, 140 and 260 lb N/ac treatments. Suboptimal N rates, especially the control, often delay crop maturation which results in wetter grain at harvest. High N rates can enhance staygreen in corn, which delays maturation and results in slightly greater grain moisture. Grain N concentration increased with increasing N rate but was statistically equal among the 180, 220, and 260 lb/ac N rates. Nitrogen removal in grain increased with increasing N rate and was statistically similar between the 220 and 260 lb N/ac rates. Grain P concentration was greatest in the control and least with 260 lb N/ac. Grain K concentration was greatest in the control and statistically similar among other N rates, while grain P₂O₅ and K₂O removal was least in the control and statistically similar among other N rates. Grain S concentrations were not affected by N rates and S removal increased with increasing N rates. Lower grain yields in 2023 resulted in less nutrient removal than previous years of this study.

Corn silage yield, silage quality and nutrient uptake in silage

Corn silage yield, silage moisture and other corn production parameters for the 2021 setup year are presented in Table 5a. Since this was a setup year the crop systems (corn for grain vs corn for silage) and cover crop treatments had not been established; therefore, the only true treatment effects would have been from the N rates applied for corn after the previous year's soybean. Silage yield at harvest moisture (wet) was numerically least with the control treatment (#1) but no significant differences among treatments were found (P value = 0.170). Wet silage yields in N fertilized treatments ranged from 22.7 to 24.9 ton/ac. Dry silage yields were less with the N control (7.47 ton/ac), however all other treatments had statistically equal yields ranging from 8.41 to 8.99 tons of dry matter per acre (TDM/ac). Silage moisture at harvest was not affected by treatments and ranged from 61.6 to 63.8%. Plant population averaged slightly less than 30,000 and was not affected by N treatments. Relative leaf chlorophyll content calculated from SPAD readings was least with the N control, intermediate with 110 lb N/ac and statistically equal with all other treatments.

Corn silage yield, silage moisture and other corn production parameters in 2022 are presented in Table 5b. An ANOVA of all 11 treatments showed silage yields, SPAD and RLC were least with the N control and generally reduced with 140 lb N/ac compared to higher N rates. Silage moisture was greatest with the N control treatment and generally driest with silage treatments with cover crops. Plant populations ranged from 32,900 to 34,800 and were greater in the corn silage crop system. Residue cover after planting ranged from 15 to 67.5%. Residue cover was greatest with corn for grain at 140 and 260 lb N/ac, slightly less with corn for grain at other N rates and was considerably less with corn silage treatments.

A split-plot statistical analysis of the main eight treatments can also be found in Table 5b. Treatment main effects (crop system and N rate for corn) did not affect any of the silage yield and corn production parameters and there were no significant interactions among main effects. However, crop system did affect residue cover after planting. Residue cover was least with Snc (16%) and Sblend (22%), intermediate with Srye (38%) and greatest with Gnc (54%). These data clearly show how the Snc system does not maintain adequate residue cover for erosion protection. The Sblend system is numerically better but still marginal, whereas Srye which was terminated in spring maintained >30% residue cover. In this strip-tillage system, the corn grain system had >50% residue cover after planting but also had numerically lower silage yields and SPAD values. The SPAD data suggested that Gnc needs more N to optimize production compared with the silage system. The added residue cover in the Gnc system likely immobilizes some soil and fertilizer N which partly explains the greater N demand. Furthermore, greater residue cover keeps the soil cooler in the spring which slows N mineralization from soil organic matter.

Corn silage yield, silage moisture and other corn production parameters in 2023 are presented in Table 5c. The ANOVA of all 11 treatments showed silage yields, SPAD and RLC were least with the control and generally reduced with 140 lb N/ac compared to higher N rates. Silage moisture was greatest with the control and 260 lb N/ac treatments and generally drier with silage treatments. Plant populations ranged from 33,800 to 34,200 and were not affected by treatments. SPAD and RLC were numerically greatest with the 260 lb N/ac treatment. However, several other treatments had statistically equal values when compared to the 260 lb N/ac. Residue cover after planting ranged from 18 to 66.5%. Residue cover was greatest with Gnc at 260, 220 and 140 lb N/ac, slightly less with Gnc at 4.6 (control) and 180 lb N/ac and was considerably less with corn silage treatments.

A split-plot statistical analysis of eight treatments in 2023 can also be found in Table 5c. When averaged across N rates, crop systems affected silage moisture and residue cover but did not affect other production parameters and there were no significant interactions. Silage moisture and residue cover were greater with the corn grain system than with corn silage. When averaged across crop systems, N rates only affected residue cover as residue cover was greater with 220 lb N/ac than with 180 lb N/ac. In 2023 all corn silage treatments had <30% residue cover, whereas, in 2022 the Srye treatments maintained >30% residue cover.

Silage quality parameters, nutrient concentrations, and nutrient removal in 2022 are presented in Table 6b. An ANOVA of all 11 treatments found crude protein (CP), starch and milk yield were least with the control and less with 140 lb N/ac compared to higher N rates. Generally, the control had greater acid detergent fiber (ADF), neutral detergent fiber (aNDF) and total tract NDF digestibility (TTNDFD). Silage P concentration was not affected by the 11 treatments and ranged from 0.194 to 0.223% (% of DM). Silage K concentration ranged from 0.586 to 0.775% and was greatest with the control but most treatments were not significantly different. Silage S concentrations were least with the control (0.055%) and 140 lb N/ac (0.058%) and greatest with the

Sblend at 220 lb N/ac treatment (0.074%). Nutrient removal in corn silage was always least with the control treatment and generally aligned with silage yield (Table 5b). With the control and 140 lb N/ac treatments excluded, nutrient removal in silage averaged 86, 136 and 12 lb/ac of P₂O₅, K₂O and S, respectively. When averaged across corn grain treatments at 180 and 220 lb N/ac (treatment #'s 2 and 3), nutrient removal in corn grain averaged 55, 38 and 9 lb/ac of P₂O₅, K₂O and S, respectively (Table 4b). These data are important when determining fertilizer rates for corn grain vs corn silage systems.

A split-plot statistical analysis of the main eight treatments for 2022 is found in Table 6b. When averaged across the crop systems, N rates for corn did not significantly affect most of the silage quality, nutrient concentration and nutrient removal parameters except for RumenS ($P=0.072$) and there were no significant interactions among main effects. RumenS was slightly greater with 180 lb N/ac than with 220 lb N/ac. When averaged across N rates, ADF was greatest with Srye while milk yield was least with Srye. The uNDF concentration was greatest with Srye, least with Snc, and intermediate with Gnc and Sblend. Gnc had slightly less P₂O₅ removal ($P=0.089$) than did silage systems (Snc and Sblend). All other parameters (CP, aNDF, TTNDFD, starch, P, K, S concentrations and K₂O and S removal) were not significantly affected by crop systems, when averaged across N rates. These data showed crop systems were more likely to affect quality parameters and nutrient concentrations and removal than were N rates. Furthermore, these data suggest that for the 2022 growing season, a 180 lb N/ac rate optimized yield, economic return, and forage quality.

Silage quality parameters, nutrient concentrations, and nutrient removal in 2023 are presented in Table 6c. An ANOVA of all 11 treatments found CP, starch and milk yield were least with the control. The control treatment had greater ADF, aNDF and RumenS. The 140 lb N/ac treatment resulted in lower concentrations for CP, ADF, aNDF and uNDF, although a few other treatments had statistically equal concentrations of these parameters when compared with 140 lb N/ac treatment. Milk yield was significantly less only with the control and the Srye treatments. Silage P and K concentrations were greatest in the control. Phosphorus concentration ranged from 0.197 to 0.222% as a % of DM, while silage K concentration ranged from 0.710 to 0.845%. Silage S concentrations were least with the control (0.055%) and 140 lb N/ac (0.062%) and greatest with the Snc and Sblend at 220 lb N/ac treatments. Nutrient removal in corn silage was always least with the control treatment and often less with the 140 lb N/ac treatment and the Gnc treatments. With the control and 140 lb N/ac treatments excluded, nutrient removal in silage averaged 65, 123 and 9.3 lb/ac of P₂O₅, K₂O and S, respectively. When averaged across corn grain N rate treatments ≥ 180 lb N/ac, nutrient removal in corn grain averaged 34, 27 and 7.0 lb/ac of P₂O₅, K₂O and S, respectively (Table 4c).

Several parameters (ADF, aNDF, uNDF, and K concentration) were considerably greater in 2023 compared with 2022. However, a few parameters (CP, starch and milk yield) were lower in 2023. These differences could be related to differences in yield between the two years or environmental conditions (more severe drought in 2023).

The split-plot statistical analysis of eight treatments for 2023 is found in Table 6c. When averaged across N rates, only uNDF, TTNDFD and milk yield were affected by crop systems. Milk yield was less with Srye than with Gnc and Snc while uNDF was greater with Srye than with Gnc and Snc. The TTNDFD concentration was greater with Gnc compared with silage treatments. When averaged across the crop systems, the 220 lb/ac N rate had greater CP and K₂O and S nutrient removal than the 180 lb/ac N rates. Although the nutrient removals were statistically significant the differences between N rates were rather small. There were no significant interactions among main effects in 2023. Most of the corn silage quality parameters were not affected by crop system or N rates in 2023.

Tile drainage and nitrate concentrations and loss (loads)

Treatment means and standard errors (SE) for tile flow, flow-weighted (FW) NO₃-N concentration and NO₃-N loss or load in 2022 are presented in Table 7a. Tile flow began on 8 April 2022 and 99.8% of all tile flow occurred in April, May and June (Fig. 1a). This late start for spring tile flow resulted from a dry growing season in 2021 that left the soil profile with a moisture deficit and a cool and relatively dry March and early April of 2022. Total annual flow averaged across treatments was 7.33 inches, which is less than normal. Total annual flow ranged from 4.8 (SE=1.13) inches with treatment #5 (Snc at 220 lb N/ac) to 9.8 (2.77) inches with treatment #3 (Gnc at 220 lb N/ac). Except for treatment #'s 3, 5 and 7, mean tile flow among treatments was relatively uniform. Flow differences among treatments and among replications within a treatment often result in data that are not normally distributed and have non constant variance among treatments; therefore, only monthly treatment means and standard errors (SE) are reported in Table 6a. When averaged across all 9 treatments, tile flow was 0.9, 3.9 and 2.5 inches in April, May and June, respectively. Heavy rain during the morning of 13 June resulted in flooding of some drainage wells/culverts which restricted flow and contaminated water samples. However, all drainage plots had returned to normal operation within 16 hours of the flooding.

In most treatments mean FW NO₃-N concentrations were least in April (ranged from 3.7 to 7.3 mg/L), increased in May (ranged from 2.9 to 9.9 mg/L) and were greatest in June (ranged from 4.1 to 13.8 mg/L, Table 7a). Averaged across months NO₃-N concentrations were least in the control (3.5 mg/L) and numerically greatest with the Snc at 220 lb N/ac treatment (10.5 mg/L). Both silage with rye cover treatments (#'s 6 & 7) had numerically lower FW NO₃-N concentrations compared with other fertilized crop system treatments. This was clear in May when NO₃-N concentrations in other crop system treatments increased. These data suggest the cereal rye cover was an effective scavenger of NO₃ thereby reducing NO₃-N concentrations in drainage.

Averaged across all fertilized treatments NO₃-N losses were 1.1, 6.0 and 7.0 lb/ac in April, May and June, respectively (Table 7a). The total annual NO₃-N loss in 2022 ranged from 6.3 (3.9) lb/ac in the control to 22.7 (7.4) lb/ac with Gnc at 220 lb N/ac. The large SE (7.4) in the Gnc at 220 lb N/ac treatment resulted from

considerable variability among replications within that treatment. Total annual NO₃-N loss in all other fertilized crop system treatments only ranged from 10.2 (3.0) to 14.8 (3.1) lb/ac.

Treatment means and standard errors (SE) for tile flow, flow-weighted (FW) NO₃-N concentration and NO₃-N loss in 2023 are presented in Table 7b. Tile flow began in about half of the plots on 22 March 2023 but during this early flow period 5 of the 36 plots had substantial flow while others had moderate to no tile flow (daily data not shown). This irregularity of flow was caused by water ponding in some areas of the 2.0-acre drainage site. The water ponding resulted from melting of a deep snowpack and snow drifts. By 26 March nearly all tiles were flowing, but two tiles still had much greater flow due to water ponding. By 28 March the ponded water had disappeared and tile flow among plots had normalized. Due to this unusual flow period (22 through 28 March), I separated this early erratic flow from the larger dataset. As shown in Fig 1b the cumulative flow graph was reset on March 28 and tile flow prior to this date was excluded from the year's cumulative data but is shown in Table 7b as "March". The last 3 days of March (29 through 31 March) were included or added to April for simplicity of reporting and statistical analysis. With flow from 22 through 28 March excluded, total annual flow averaged across treatments was 9.56 inches and 99.7% of all flow occurred in April and May (Fig. 1b). Total annual flow ranged from 6.1 (1.56) inches with treatment #5 (Snc at 220 lb N/ac) to 11.8 (3.12) inches with treatment #7 (Srye at 220 lb N/ac). Except for treatment #'s 5 and 7, mean tile flow among treatments was relatively uniform. When averaged across all 9 treatments, tile flow was 2.6, 4.2 and 5.3 inches in March, April and May, respectively. Large rainfall events occurred on 31 Mar through 1 April and 6 through 14 May (Fig 1b), nearly 6 inches of rainfall was recorded in these 9 days in May. These large events were responsible for most of the annual total tile flow. Some tiles (drainage wells) flooded on 14 May. However, all drainage plots had returned to normal operation within 20 hours of the flooding.

In 2023 monthly mean FW NO₃-N concentrations were numerically greater in May (ranged from 2.1 to 15.7 mg/L) than in March or April (April ranged from 1.9 to 10.8 mg/L, Table 7b). When averaged across April and May, FW NO₃-N concentrations were least in the control (2.1 mg/L) and numerically greatest with the Snc at 220 lb N/ac and Sblend treatments (13.5 mg/L). Nitrate-N concentrations in the Srye treatments (#'s 6 & 7) were nearly equal to other silage crop system treatments. However, the grain system treatments had lower FW NO₃-N concentrations than the silage systems.

When averaged across all fertilized treatments, NO₃-N losses were 3.8, 9.0 and 15.7 lb/ac in March, April and May, respectively (Table 7b). Total annual NO₃-N loss in 2023 ranged from 5.0 (3.0) lb/ac in the control to 33.2 (6.9) lb/ac with Sblend at 220 lb N/ac. The Snc at 220 lb N/ac treatment had lower total annual NO₃-N loss compared to other fertilized treatments. This was due to this treatment having lower total annual flow, only 6.1 inches and it had the least annual flow in 2022 too.

Crop system, cover crop and N rate treatment effects on annual and three-month interval FW NO₃-N concentration and annual flow-adjusted NO₃-N losses in 2022 are presented in Table 8a. Due to 98.8% of the entire annual flow occurring during the April-June period, the April-June and annual average FW NO₃-N concentration data and interpretation are almost identical. Therefore, only the April-June concentration data will be discussed here. An ANOVA of all 9 treatments showed FW NO₃-N concentration was least with the control (2.7 mg/L) followed by the Srye at 180 lb N/ac treatment (5.9 mg/L). The Srye at 180 lb N/ac treatment #6 had lower FW NO₃-N concentrations than treatment #'s 3, 4, 5 and 9. The Snc at 220 lb N/ac treatment had greater FW NO₃-N concentration than all other treatments except the Gnc at 220 lb N/ac. An ANOVA of the 8 fertilized treatments found when averaged across N rates, FW NO₃-N concentrations in April-June and the annual average were greater with Snc than with Srye and Sblend. Srye had 36% lower NO₃-N concentrations than did Snc. When averaged across crop systems, 220 lb N/ac had greater flow-weighted NO₃-N concentration than 180 lb N/ac.

Nitrate-N loss or load is calculated by multiplying tile flow (gal/ac/day) by a constant (8.34 lb/gal) and by the NO₃-N concentration in the sample for that period (a few days during heavy flow or a week or more during light tile flow). Flow-weighted NO₃-N concentrations are calculated by summing the NO₃-N loss for a period of days (3 months or the annual total) and dividing by the sum of the flow and the constant for the same period. Flow-adjusted NO₃-N loss is calculated by dividing the annual sum of nitrate loss by the sum of the total annual flow (Eq. 1). Flow-adjusted NO₃-N loss data are normalized for variability in tile flow among individual plots and treatment means. This adjustment or correction minimizes variation and usually allows these data to be analyzed with ANOVA without violating the normality and constant variance assumptions.

$$\text{Equation 1} \quad \text{total annual nitrate lost} \div \text{total annual flow} = \text{flow adjusted loss}$$

An ANOVA of all 9 treatments showed April-June and the annual total NO₃-N loss was least with the control (4.3 lb/ac) and greatest with Gnc at 220 lb N/ac treatment (22.7 lb/ac, Table 8a). All other treatments had statistically equal April-June and the annual total NO₃-N loss. An ANOVA of the 8 fertilized treatments found no significant differences (P value = 0.442) in NO₃-N loss in April-June and the annual average, when averaged across N rates. The lack of significant differences in NO₃-N loss is not uncommon in drainage research studies. The variability in flow among plots and replications increases the mean squared error term which reduces the *F* statistic in ANOVA. When averaged across crop systems, 220 lb N/ac had 33% greater NO₃-N loss than 180 lb N/ac. The 180 lb/ac N rate falls within the Profitable N Rate Range of 159 to 189 lb/ac for corn after corn in Minnesota (0.10 price ratio) from the [Corn N Rate Calculator](#). The 180 and 220 lb/ac N rates had statistically equal corn silage and grain yields in 2022 (Tables 4b and 5b). However, corn grain yields were greater with 260 lb N/ac but that treatment did not have tile drainage monitoring.

An ANOVA of all 9 treatments showed flow-adjusted NO₃-N loss was least with the control (0.62 lb/inch) followed by the Srye at 180 lb N/ac treatment (1.35 lb/inch), and greatest with Snc at 220 lb N/ac (2.82 lb/inch, Table 8a). An ANOVA of the 8 fertilized treatments found flow-adjusted NO₃-N loss was reduced significantly with Srye and Sblend compared with Snc, when averaged across N rates. Srye had 36% lower flow-adjusted NO₃-N loss than Snc. When averaged across crop systems, 220 lb N/ac had greater flow-adjusted NO₃-N loss than 180 lb N/ac.

Crop system, cover crop and N rate treatment effects on annual and three-month interval FW NO₃-N concentration and annual flow-adjusted NO₃-N losses in 2023 are presented in Table 8b. An ANOVA of all 9 treatments showed FW NO₃-N concentration was least with the control and less with Gnc at 180 lb N/ac for both (January-March and April-June) 3-month intervals. Since the erratic late March flow data was removed from the annual total and very little flow occurred after 30 June, the total annual flow is mostly from the April-June interval. An ANOVA of the 8 fertilized treatments found FW NO₃-N concentrations for the April-June interval and the annual average were less with the corn grain system (Gnc) than with silage systems (Snc, Srye and Sblend). In 2022 Srye had lower FW NO₃-N concentrations than Gnc. The rye cover crop was not effective at sequestering N from the soil in 2023, likely due to poor rye growth in the fall of 2022 and spring of 2023. When averaged across crop systems, 220 lb N/ac had greater FW NO₃-N concentrations than 180 lb N/ac for both 3-month intervals and the annual average.

An ANOVA of all 9 treatments showed the April-June interval and annual total NO₃-N loss was least in the control (5.0 lb/ac) and numerically greatest with Sblend at 220 lb N/ac (33.2 lb/ac) and Srye at 220 lb N/ac (31.5 lb/ac) treatments (Table 8b). Nitrate-N loss was less with the Gnc at 180 lb N/ac treatment than with almost all the other fertilized treatments. The Snc at 220 lb N/ac treatment had less NO₃-N loss than Srye and Sblend at 220 lb N/ac treatments, but like 2022 it had less tile flow. An ANOVA of the 8 fertilized treatments found no significant differences in NO₃-N loss for the April-June interval (P value = 0.120) and the annual average (P value = 0.119), when averaged across N rates. When averaged across crop systems, 220 lb N/ac had numerically (16%) greater NO₃-N loss than 180 lb N/ac but it was not statistically significant (P value = 0.190). Like 2022, the 180 and 220 lb/ac N rates had statistically equal corn silage and grain yields in 2023 (Tables 4c and 5c). However, corn grain yields were greater with 260 lb N/ac than with 180 lb N/ac.

An ANOVA of all 9 treatments found flow-adjusted NO₃-N loss in 2023 was least with the control (0.47 lb/inch) followed by the Gnc at 180 lb N/ac, treatment #2 (1.35 lb/inch), and greatest with Snc at 220 lb N/ac (3.08 lb/inch) and Sblend at 220 lb N/ac (3.07 lb/inch, Table 8b). An ANOVA of the 8 fertilized treatments found flow-adjusted NO₃-N loss was significantly less with the grain corn system (Gnc) than with silage systems (Snc, Srye and Sblend), when averaged across N rates. Gnc had 38% less flow-adjusted NO₃-N loss than Snc. When averaged across crop systems, 220 lb N/ac had greater flow-adjusted NO₃-N loss than 180 lb N/ac.

Soil nitrogen

The effects of N rates for corn in 2021 on total residual soil NO₃-N (RSN) and NH₄-N remaining in the soil profile are shown in Figure 2. Nitrate-N increased at the 200 lb N/ac rate compared with other N rates in the corn grain crop system. However, N rates had little effect on NH₄-N and NH₄-N was quite low (<18 lb/ac) in the 0- to 90-cm profile. The effects of cover crops and N rates on residual soil NO₃-N and NH₄-N remaining in the soil profile in the corn silage system is shown in Figure 3. Nitrate was least, only 16.6 lb/ac, with the rye cover at 140 lb N/ac and was greater with the annual blend cover. Greater N rates resulted in greater RSN in the soil profile with rye and blend covers. Oddly, the Snc at 140 lb N/ac rate had greater RSN than Snc at 170 lb N/ac. These data show the potential of the Srye system to scavenge RSN from the soil profile and thereby potentially reduce NO₃-N concentrations and losses in tile drainage water. Whereas, the Sblend system had similar RSN as the Snc system in early May of 2022.

The effects of crop system and N rates for corn in 2022 on soil inorganic N in the 0- to 30-cm depth were measured on 10 November 2022 (Table 10). Nitrate-N concentrations ranged from 2.5 mg/kg in the control to 15.9 mg/kg with Srye at 220 lb N/ac. When averaged across N rates, soil NO₃-N concentrations were numerically less with Gnc, than with corn silage systems (Snc, Srye and Sblend). However, these differences were not statistically significant (P value = 0.148). When averaged across crop systems, soil NO₃-N concentrations were greater with 220 lb N/ac (11.8 mg/kg) than with 180 lb N/ac (7.6 mg/kg). No differences in soil NO₃-N concentrations were observed among the silage crop systems which means there was no or very little effect of cover crops on soil nitrate in the 0- to 30-cm depth in the fall of 2022. Due to a very dry September and early October cover crop growth was extremely small, so a cover crops ability to scavenge NO₃-N would have been minimal. The numeric differences among N rates, including the control (4.6 lb N/ac), in the corn grain system were small, suggesting the 180 and 220 lb N/ac rates left very little N in the soil profile after harvest. As they were only slightly greater than the control treatment. Crop systems and N rates for corn in 2022 had no significant effects on NH₄-N concentrations in the fall of 2022.

The effects of crop system and N rates for corn in 2023 on soil inorganic N in the 0- to 30-cm depth were measured on 4 November 2023 (Table 10). Nitrate-N concentrations ranged from 1.2 mg/kg in the control to 25.2 mg/kg with Snc at 220 lb N/ac. When averaged across N rates, soil NO₃-N concentrations were less with Gnc (5.2 mg/kg), than with Snc (19.8 mg/kg) and Srye (15.7 mg/kg), while the Sblend treatment was intermediate (13.0 mg/kg). This 2023 data showed that in the late fall silage corn systems even with a cover crop are at greater risk for nitrate loss than corn grain systems. Furthermore, the yield data showed grain yields increased up to 260 lb N/ac; whereas, silage yields were not different between the 180 and 220 lb/ac N rates. These yield data suggest the 180 lb N/ac rate is sufficient in corn silage systems but likely needs to be greater than the current Profitable N Rate Range of 159 to 189 lb/ac for corn grain systems. When averaged across crop systems, soil NO₃-N concentrations were not significantly affected by N rates. No significant differences in main effects (crop systems and N rates for corn) were found for soil NH₄-N concentrations in

2023. However, a significant interaction between main effects for NH₄-N concentrations was found. Soil NH₄-N concentrations were less with Sblend at 220 lb N/ac than with Sblend at 180 lb N/ac, while with other crop systems NH₄-N concentrations were numerically greater with 220 lb N/ac than with 180 lb N/ac. The authors have no explanation for this finding however, the same trend was observed in NO₃-N concentrations in 2023.

Soil physical and biological parameters

A preliminary analysis of soil physical parameters (bulk density, BD) was conducted and presented in last year's annual report. Some biological parameters [POXC and CO₂ burst (min-C)] from the 2023 sampling have only recently been analyzed and were not included in this report. Spring soil samples from all years (2022 through 2024) will be shipped to the lab and analyzed for total organic carbon in June of 2024. Analyzing all years together improves quality control of the data. Analysis and interpretation of the 2022 and 2023 spring soil data is on-going; therefore, these data were not included in this annual report but will be presented and discussed in later reports and outreach activities.

Results Summary

A research study initiated in 2021 continued in 2023 the objectives were to quantify the effects and interactions of cover crops and N rates on corn production, nitrate-N concentration and loss in tile drainage, soil N and soil health parameters. Cover crops (cereal rye and annual blend) were drilled after corn silage harvest each fall. Cover crop biomass yields (dry matter yield) on 5 November 2021 were numerically greater with rye (296 lb) than with blend (255 lb, *P* value = 0.114). In spring prior to termination, rye biomass was nearly equal to the previous fall. An extraordinary dry fall resulted in very poor growth of cover crops in 2022, biomass yields of rye on 8 May 2023 averaged <30 lb/ac. Cover crop biomass yield on 23 October 2023 averaged 92 and 229 lb/ac for rye and blend, respectively.

Corn grain yields ranged from 111 bu/ac in the control (4.6 lb N/ac) to 167 bu/ac with 170 lb N/ac for corn following soybean in 2021. An ANOVA analysis showed yields were statistically equal among the 140, 170 and 200 lb N/ac treatments. Corn grain yields ranged from 60 and 48 bu/ac in the control to 218 and 163 bu/ac with 260 lb N/ac in 2022 and 2023, respectively. A cool and wet spring delayed planting until 22 May which was followed by a summer drought. Both negatively affected corn growth and production in 2023. Silage yields in 2021 were only slightly less with the control than other N rates. Silage yields in 2022 and 2023 were not significantly affected by crop systems (corn grain production or corn silage with or without cover crops) or N rates (180 vs 220 lb/ac). In 2022 and 2023 corn grain yields generally increased with increasing N rate while silage yields were less affected by N rates except for the control (4.6 lb N/ac). This corn grain and silage yield response to N rate suggests a greater N rate is needed for grain production than with silage production. Cooler soils due to greater residue cover in the corn grain system could reduce N mineralization of SOM and increase N immobilization. Corn silage quality was only minimally affected by the crop system and N rate treatments. However, low N rates, especially the control often affected silage quality. The control caused some quality

parameters to increase and others to decrease. Nutrient uptake in silage was much less in the control due to reduced yield.

Annual tile drainage averaged across treatments was 7.3 and 9.6 inches in 2022 and 2023, respectively. Nearly all drainage occurred during the months of April, May and June in 2022 and March, April and May in 2023, which is unusual especially in the last 20 years. Flow-weighted NO₃-N concentrations and flow-adjusted losses in tile drainage water averaged 20% greater with 220 lb N/ac than with 180 lb N/ac. In 2022, annual FW NO₃-N concentrations and flow-adjusted losses were reduced by 36 and 21% by cereal rye and annual blend (cover crops), respectively. However, cover crops had no effect on FW NO₃-N concentrations and flow-adjusted losses in 2023. Cover crops had sufficient growth in the fall of 2021 and spring of 2022 but minimal growth in the fall of 2022 and spring of 2023. Cover crop growth (biomass yield) clearly impacts NO₃-N concentrations and losses in tile drainage. In 2023, FW NO₃-N concentrations were less with the corn grain system (Gnc, 8.1 mg/L) compared with corn silage systems (12.1 to 13.0 mg/L). For comparison FW NO₃-N concentrations in the control treatment which received 4.6 lb N/ac were 2.7 and 2.1 mg/L in 2022 and 2023, respectively. Annual flow-adjusted NO₃-N losses ranged from 0.82 and 0.47 lb/inch with the control to 2.82 and 3.08 lb/inch with Snc at 220 lb N/ac in 2022 and 2023, respectively.

Inorganic N concentrations from soil samples taken in the fall (late October or early November) each year showed NO₃-N concentrations at a 0- to 30-cm depth were less with the corn grain crop system (Gnc) than with silage systems. Cover crops seeded after silage harvest did not significantly reduce soil NO₃-N concentrations in these fall samples. However, Sblend had numerically less soil NO₃-N in 2023. Nitrate-N concentrations were greater with 220 lb N/ac than with 180 lb N/ac in 2022. Soil NH₄-N concentrations were not affected by crop system or N rates.

Outreach and Extension Activities

Preliminary findings of this research project were presented at meetings on 10 August 2022 (Soil Health Nexus), 15 November 2022 (MASWCD Area V1 Meeting), 6 December 2022 (AFREC Research Update), 20 June 2023 (SROC Agronomy Tour), and 12 December 2023 (AFREC Research Update).

Acknowledgement

This work was supported by the Agricultural Fertilizer Research and Education Council (AFREC). The author greatly appreciates their support. The author also thanks co-investigator (Dr. Anna Cates) for her input and expertise and Kyle Holling for tile water monitoring and sample collection.

References

- Badger, S. D. E. Kaiser, and M. S. Wells. 2017. Nitrogen Availability and Corn Production in Minnesota Following Cover Crops. ASA CSSA SSSA Annual Meeting. Oct. 22-25 Tampa, FL. Online: <https://scisoc.confex.com/scisoc/2017am/webprogram/Paper107347.html>
- Dinnes, D.L., D.L. Karlen, D.B. Jaynes, T.C. Kaspar, J.L. Hatfield, T.S. Covlin and C.A. Cambardella. 2002. Nitrogen management strategies to reduce nitrate leaching in tile-drained Midwestern Soils. *Agron. J.* 94:153-171.
- Kaspar, T.C., D.B. Jaynes, T. Parkin, and T. Moorman. 2007. Rye Cover Crop and Gamagrass Strip effects on nitrate concentration and load in tile drainage. <https://doi.org/10.2134/jeq2006.0468> *J. Environ. Qual.* 36(5):1503-11.
- Oldfield, E. E., Bradford, M. A., Augarten, A. J., Cooley, E. T., Radatz, A. M., Radatz, T., & Ruark, M. D. 2021. Positive associations of soil organic matter and crop yields across a regional network of working farms. <https://doi.org/10.1002/saj2.20349> *Soil Science Society of America Journal.* 86(2):384-397.
- Randall, G.W., and D.J. Mulla. 2001. Nitrate nitrogen in surface waters as influenced by climatic conditions and agricultural practices. *J. Environ. Qual.* 30:337–344.
- Snyder C.S., and T.W. Bruulsema. 2007. IPNI Publ. No. 07076. Norcross, GA., U.S.A. pp. 4.
- Strock, J.S., P.M. Porter, and M.P. Russelle. 2004. Cover crop to reduce nitrate loss through subsurface drainage in the northern Corn Belt. *J. Environ. Qual.* 33:1010-1016.

Table 1. Experimental methods and dates of completion (2021 was a setup year).

Experimental method or procedure	Study year			
	2021	2022	2023	2024
	----- Month / day -----			
Previous crop	Soybean	Corn/silage	Corn/silage	Corn/silage
Collect spring soil samples	--	5/6	5/1	
Collect cover crop biomass samples	NA	4/27	5/8	
Cereal rye terminated with herbicide	NA	4/27, 5/18	5/17	
Corn planting & starter fertilizer	5/7, 5/26	5/19	5/22	
Residue measurements	NA	6/1	6/5	
Pre-emerge herbicide application	5/11	5/27	5/17	
V2 N application	6/14	6/9	6/8	
Corn plant stand counts	6/24	6/23	6/13	
Post-emerge herbicide application	7/1	6/24	6/15	
Take plot notes at V8 and R3	7/8 (V9)	7/6, 8/19	7/6, 8/17	
SPAD measurements at R1	8/2	8/1	8/1	
Hand harvest silage plots	9/9	9/13	9/6	
Clean off remaining corn silage	9/14	9/16	9/7	
Seed cover crops	9/14	9/17	9/13	
Combine harvest corn grain	10/14	10/14	10/21	
Clean off remaining corn grain plots	10/16	10/21	11/1	
Collect cover crop biomass samples	10/27	10/31	10/23	
Collect corn stover (residue) samples	--	11/28	11/3	
Shred corn stalks on grain plots	11/3	10/31	11/4	
Strip till and apply P-K-S fertilizers	11/5	11/1	11/5	
Collect fall soil samples	11/24	11/10	11/4	

Table 2. Monthly total precipitation, mean air temperature, and growing degree units (GDU, base 50/86) as compared to 30-year normal values at Waseca.

Month	Year	Precipitation		Mean Air Temp.		GDUs	
		Observed	Normal [†]	Observed	Normal [†]	Observed	Normal [†]
		----- inches -----		----- °F -----			
Jan	2021	1.04	1.27	19.2	13.2	-	-
Feb	2021	0.67	1.20	7.5	17.5	-	-
Mar	2021	2.41	2.25	38.2	30.7	-	-
Apr	2021	0.62	3.30	45.9	45.1	-	-
May	2021	2.66	4.47	58.1	57.9	344	320
Jun	2021	2.00	5.38	74.0	68.4	674	548
Jul	2021	2.73	4.93	71.9	71.4	666	652
Aug	2021	4.82	4.82	70.8	69.0	645	591
Sep	2021	1.92	4.12	63.7	61.6	452	374
Oct	2021	2.98	2.77	53.8	47.8	199	24
Nov	2021	0.64	1.89	34.9	32.9	-	-
Dec	2021	1.69	1.50	23.4	19.8	-	-
May-Sep	Total	14.13	23.72	66.8	65.7	2,629	2484
Annual	Total	24.18	37.90	46.8	44.6	2,979	2509
Jan	2022	1.13	1.27	6.5	13.2	-	-
Feb	2022	0.69	1.20	11.2	17.5	-	-
Mar	2022	1.62	2.25	30.2	30.7	-	-
Apr	2022	3.75	3.30	38.7	45.1	-	-
May	2022	4.74	4.47	59.1	57.9	336	320
Jun	2022	4.36	5.38	70.3	68.4	594	548
Jul	2022	4.60	4.93	72.5	71.4	691	652
Aug	2022	5.50	4.82	69.3	69.0	596	591
Sep	2022	0.78	4.12	62.8	61.6	412	374
Oct	2022	0.36	2.77	48.6	47.8	0	24
Nov	2022	1.84	1.89	33.2	32.9	-	-
Dec	2022	2.03	1.50	14.8	19.8	-	-
May-Sep	Total	19.98	23.72	66.8	65.7	2,629	2,484
Annual	Total	31.40	37.90	43.1	44.6	2,629	2,509
Jan	2023	2.43	1.27	17.1	13.2	-	-
Feb	2023	2.51	1.20	15.9	17.5	-	-
Mar	2023	2.35	2.25	26.2	30.7	-	-
Apr	2023	3.66	3.30	44.4	45.1	-	-
May	2023	6.47	4.47	61.5	57.9	401	320
Jun	2023	1.56	5.38	72.6	68.4	661	548
Jul	2023	1.62	4.93	69.7	71.4	602	652
Aug	2023	3.28	4.82	71.2	69.0	642	591
Sep	2023	2.26	4.12	66.8	61.6	500	374
Oct	2023	3.78	2.77	50.3	47.8	127	24
Nov	2023	0.09	1.89	35.4	32.9	-	-
Dec	2023	1.77	1.50	32.1	19.8	-	-
May-Sep	Total	15.19	23.72	68.4	65.7	2,806	2484
Annual	Total	31.78	37.90	46.9	44.6	2,933	2509

[†] 30-Yr normal, 1991-2020.

Table 3a. Cover crop dry matter yield, nutrient concentration and nutrient uptake as affected by cover crop species and N rates for corn.

Trt	Treatments		Cover crop biomass on 5 Nov 2021						Cover crop biomass on 27 Apr 2022					
	Cover crop	N rate	Yield	N conc.	C conc.	C:N ratio	N uptake	C uptake	Yield	N conc.	C conc.	C:N ratio	N uptake	C uptake
#		lb/ac	lb/ac	----- % -----			----- lb/ac -----		lb/ac	----- % -----		----- lb/ac -----		
6	Cereal rye	140	282	3.21	39.5	12.3	9.0	111	291	3.03	38.3	12.6	8.8	112
7	Cereal rye	170	311	3.89	41.1	10.6	12.1	128	301	3.11	38.4	12.3	9.4	116
8	Annual blend	140	254	3.89	41.4	10.8	10.0	105						
9	Annual blend	170	257	4.21	40.7	9.7	10.8	105						

Stats for RCB Design with a split-plot arrangement

Cover crop (main plot)

Cereal rye	296 A [†]	3.55 B	41.4 A	11.4 A	10.6 A	119 A
Annual blend	255 A	4.05 A	41.1 A	10.2 B	10.4 A	105 A
P > F:	0.114	0.042	0.401	0.059	0.868	0.134

N rate for corn in 2021 (sub-plot)

140	268 A	3.55 B	41.5 A	11.5 A	9.5 B	108 A	291 A	3.03 A	38.3 A	12.6 A	8.8 A	112 A
170	284 A	4.05 A	40.9 B	10.1 B	11.4 A	116 A	301 A	3.11 A	38.4 A	12.3 B	9.4 A	116 A
P > F:	0.418	0.014	0.067	0.009	0.072	0.300	0.871	0.227	0.982	0.092	0.738	0.856

Interaction (cover crop × N rate)

P > F:	0.501	0.276	0.835	0.437	0.235	0.268
--------	-------	-------	-------	-------	-------	-------

† Numbers followed by different letters are significantly different at $\alpha = 0.10$ level. Capital letters signify differences in main effects and small letters are differences due to interaction between main effects.

Table 3b. Cover crop dry matter yield, nutrient concentration and nutrient uptake as affected by cover crop species and N rates for corn.

Trt	Treatments		Cover crop biomass on fall of 2022					Cover crop biomass on 8 May 2023							
	Cover crop	N rate	Yield	N conc.	C conc.	C:N ratio	N uptake	C uptake	Yield	N conc.	C conc.	C:N ratio	N uptake	C uptake	
#		lb/ac	lb/ac	----- % -----			----- lb/ac -----		lb/ac	----- % -----		----- lb/ac -----			
6	Cereal rye	180							31	4.14	40.1	9.7	1.3	13	
7	Cereal rye	220	no data, not enough growth							24	4.04	40.1	9.9	1.0	10
8	Annual blend	180													
9	Annual blend	220													

Stats for RCB Design with a split-plot arrangement

Cover crop (main plot)

Cereal rye
Annual blend
P > F:

N rate for corn in 2022 (sub-plot)

180	31 A	4.14 A	40.1 A	9.7 A	1.3 A	13 A
220	24 A	4.04 A	40.1 A	9.9 A	1.0 A	10 A
P > F:	0.636	0.504	0.873	0.606	0.609	0.638

Interaction (cover crop × N rate)

P > F:

† Numbers followed by different letters are significantly different at $\alpha = 0.10$ level. Capital letters signify differences in main effects and small letters are differences due to interaction between main effects.

Table 3c. Cover crop dry matter yield, nutrient concentration and nutrient uptake as affected by cover crop specie and N rates for corn.

Trt	Treatments		Cover crop biomass on 23 Oct 2023						Cover crop biomass on ___ May 2024					
	Cover crop	N rate	Yield	N conc.	C conc.	C:N ratio	N uptake	C uptake	Yield	N conc.	C conc.	C:N ratio	N uptake	C uptake
#		lb/ac	lb/ac	----- % -----			----- lb/ac -----		lb/ac	----- % -----		----- lb/ac -----		
6	Cereal rye	180	93	4.46 a	42.3	9.5 ab	4.2	39						
7	Cereal rye	220	91	4.25 b	41.5	9.8 a	3.9	38						
8	Annual blend	180	231	4.59 a	42.5	9.3 ab	10.8	98						
9	Annual blend	220	228	4.76 a	42.6	9.0 b	10.9	97						

Stats for RCB Design with a split-plot arrangement

Cover crop (main plot)

Cereal rye	92 B	4.35 A	41.9 A	9.6 A	4.0 B	38 B
Annual blend	229 A	4.68 A	42.5 A	9.1 A	10.8 A	97 A
P > F:	0.017	0.125	0.337	0.157	0.023	0.016

N rate for corn in 2022 (sub-plot)

180	162 A	4.53 A	42.4 A	9.4 A	7.5 A	69 A
220	159 A	4.50 A	42.0 A	9.4 A	7.4 A	67 A
P > F:	0.899	0.750	0.440	0.978	0.920	0.891

Interaction (cover crop × N rate)

P > F:	0.974	0.044	0.322	0.058	0.857	0.989
--------	-------	-------	-------	-------	-------	-------

† Numbers followed by different letters are significantly different at $\alpha = 0.10$ level. Capital letters signify differences in main effects and small letters are differences due to interaction between main effects.

Table 4a. Corn grain yield and moisture, nutrient concentration and uptake in 2021.

Trt #	Treatments			Yield bu/ac	Moisture -- % --	Grain nutrient concentration				Grain nutrient removal			
	Corn for #	Cover #	'21 N rate lb/ac			N	P	K	S	N	P ₂ O ₅	K ₂ O	S
1	Grain	None	4.6	111 c	20.0 b								
10	Grain	None	110	147 b	21.1 a	no data in 2021				no data in 2021			
2	Grain	None	140	160 ab	20.5 ab								
3	Grain	None	170	167 a	20.8 ab								
11	Grain	None	200	166 a	21.2 a								

Stats for RCB Design with a split-plot arrangement

Nitrogen rate for corn in 2021

P > F:	0.002	0.092
LSD 0.10:	18	0.8

Table 4b. Corn grain yield and moisture, nutrient concentration and uptake in 2022.

Trt #	Treatments			Yield bu/ac	Moisture -- % --	Grain nutrient concentration				Grain nutrient removal			
	Corn for #	Cover #	'22 N rate lb/ac			N	P	K	S	N	P ₂ O ₅	K ₂ O	S
1	Grain	None	4.6	60 c	24.1 bc	0.90 b	0.336 a	0.378 a	0.093 b	26 d	21.9 b	12.9 b	2.7 c
10	Grain	None	140	189 b	25.6 ab	1.07 a	0.243 bc	0.318 b	0.094 ab	96 c	50.4 a	34.3 a	8.4 b
2	Grain	None	180	194 b	22.7 c	1.16 a	0.264 b	0.348 ab	0.098 a	107 bc	55.5 a	38.3 a	9.0 a
3	Grain	None	220	201 b	23.9 c	1.17 a	0.247 bc	0.335 ab	0.095 ab	110 ab	54.5 a	38.5 a	9.0 a
11	Grain	None	260	218 a	26.2 a	1.18 a	0.196 c	0.273 c	0.086 c	121 a	46.4 a	33.7 a	8.9 a

Stats for RCB Design with a split-plot arrangement

Nitrogen rate for corn in 2022

P > F:	<0.001	0.020	0.003	0.017	0.013	0.014	<0.001	0.012	<0.001	<0.001
LSD 0.10:	15	1.7	0.11	0.059	0.044	0.005	12	15.3	6.9	0.5

Table 4c. Corn grain yield and moisture, nutrient concentration and uptake in 2023.

Trt #	Treatments			Yield bu/ac	Moisture -- % --	Grain nutrient concentration ----- % -----				Grain nutrient removal ----- lb/ac -----			
	Corn for #	Cover #	'23 N rate lb/ac			N	P	K	S	N	P ₂ O ₅	K ₂ O	S
1	Grain	None	4.6	48 d	21.7 a	0.85 c	0.335 a	0.368 a	0.091	19.0 d	17.2 b	9.9 b	2.0 d
10	Grain	None	140	137 c	21.4 ab	1.00 b	0.247 b	0.320 b	0.092	64.3 c	37.2 a	25.1 a	5.9 c
2	Grain	None	180	150 b	19.4 c	1.02 ab	0.215 bc	0.313 b	0.094	72.0 b	35.1 a	26.7 a	6.7 b
3	Grain	None	220	158 ab	20.4 bc	1.07 ab	0.202 bc	0.303 b	0.094	79.6 a	34.9 a	27.2 a	7.0 ab
11	Grain	None	260	163 a	21.2 ab	1.09 a	0.185 c	0.290 b	0.096	84.1 a	32.8 a	26.9 a	7.4 a

Stats for RCB Design with a split-plot arrangement

Nitrogen rate for corn in 2023

P > F:	<0.001	0.026	0.001	0.003	0.056	0.744	<0.001	0.041	<0.001	<0.001
LSD 0.10:	11	1.2	0.08	0.055	0.042	NS	5.8	10.9	4.9	0.4

Table 5a. Corn silage yield, plant population, leaf chlorophyll at R1 and residue cover after planting in 2021.

Trt #	Treatments			Corn silage yield		Silage	Plant pop. pl/ac × 10 ³	SPAD	RLC %
	Corn for Cover	'21 N rate lb/ac		Wet tons ton/ac	Dry tons ton/ac	Moisture %			
1	Grain	NA	4.6	20.4	7.47 b	63.2	29.5	45.0 d	78.2 d
2	Grain	NA	140	23.2	8.58 a	63.0	29.1	55.7 ab	97.0 abc
3	Grain	NA	170	23.1	8.62 a	62.6	29.1	55.4 abc	96.4 abc
4	Silage	NA	140	24.3	8.86 a	63.4	29.6	55.1 abc	96.0 abc
5	Silage	NA	170	24.9	8.99 a	63.8	29.6	55.8 ab	97.2 ab
6	Silage	NA	140	22.8	8.57 a	62.4	29.6	55.9 ab	97.3 ab
7	Silage	NA	170	24.4	8.93 a	63.4	29.6	55.4 abc	96.4 abc
8	Silage	NA	140	22.9	8.45 a	63.1	29.3	54.6 bc	95.2 bc
9	Silage	NA	170	23.5	8.58 a	63.5	29.6	55.4 abc	96.4 abc
10	Grain	NA	110	22.7	8.41 a	63.4	no data	53.9 c	94.3 c
11	Grain	NA	200	23.3	8.93 a	61.6	no data	56.3 a	98.0 a

Stats for RCB Design all 11 treatments

Treatment							
P > F:		0.170	0.039	0.397	0.172	<0.001	<0.001
LSD 0.10:		NS	0.62	NS	NS	1.5	2.7

Table 5b. Corn silage yield, plant population, leaf chlorophyll at R1 and residue cover after planting in 2022.

Trt #	Treatments			Corn silage yield		Silage	Plant pop. pl/ac × 10 ³	SPAD	RLC %	Residue Cover %
	Corn for	Cover	'22 N rate lb/ac	Wet tons ton/ac	Dry tons ton/ac	Moisture %				
1	Grain	None	4.6	11.4 d	3.63 d	68.3 a	33.6 bcd	36.2 e	61.9 e	50.0 b
2	Grain	None	180	23.3 bc	8.17 bc	65.0 bc	33.2 cd	55.8 cd	95.5 cd	55.5 b
3	Grain	None	220	23.3 bc	8.45 abc	63.8 cde	33.5 bcd	55.9 cd	95.6 cd	52.0 b
4	Silage	None	180	25.6 a	9.03 a	64.8 bc	34.8 a	57.1 abc	97.6 abc	15.0 e
5	Silage	None	220	24.9 ab	8.89 ab	64.3 cde	33.8 bcd	57.5 ab	98.3 ab	17.0 e
6	Silage	Rye	180	25.1 ab	9.11 a	63.7 cde	34.4 ab	56.9 abcd	97.3 abcd	35.5 c
7	Silage	Rye	220	24.2 abc	8.95 a	63.1 e	34.2 abc	56.2 bcd	96.2 bcd	40.5 c
8	Silage	Blend	180	24.8 ab	9.09 a	63.3 de	34.0 abc	57.5 ab	98.4 ab	18.5 de
9	Silage	Blend	220	25.0 ab	9.07 a	63.7 cde	34.8 a	58.2 a	99.5 a	25.0 d
10	Grain	None	140	22.3 c	7.85 c	64.7 bcd	33.3 cd	55.5 d	94.9 d	67.5 a
11	Grain	None	260	25.8 a	8.82 ab	65.9 b	32.9 d	56.5 bcd	96.6 bcd	64.5 a

Stats for RCB Design all 11 treatments

Treatment

P > F:	<0.001	<0.001	<0.001	0.025	<0.001	<0.001	<0.001
LSD 0.10:	2.0	0.74	1.5	1.0	1.4	2.5	7.8

Stats for RCB Design with a split-plot arrangement for treatments 2 through 9

Crop and cover crop system

Grain corn, no cover crop	23.3 A	8.31 A	64.4 A	33.4 A	55.8 A	95.5 A	53.8 A
Silage corn, no cover crop	25.3 A	8.96 A	64.5 A	34.3 A	57.3 A	97.9 A	16.0 C
Silage corn, cereal rye	24.7 A	9.03 A	63.4 A	34.3 A	56.5 A	96.8 A	38.0 B
Silage corn, annual blend	24.9 A	9.08 A	63.5 A	34.4 A	57.8 A	98.9 A	21.8 C
P > F:	0.289	0.235	0.313	0.114	0.132	0.170	<0.001

Nitrogen rate for corn in 2022

180 lb/ac	24.7 A	8.85 A	64.2 A	34.1 A	56.8 A	97.2 A	31.1 A
220 lb/ac	24.4 A	8.84 A	63.7 A	34.1 A	56.9 A	97.4 A	33.6 A
P > F:	0.411	0.936	0.253	0.962	0.714	0.753	0.288

Interaction (crop system × N rate)

P > F:	0.777	0.751	0.529	0.131	0.521	0.517	0.441
--------	-------	-------	-------	-------	-------	-------	-------

Table 5c. Corn silage yield, plant population, leaf chlorophyll at R1 and residue cover after planting in 2023.

Trt #	Treatments			Corn silage yield		Silage	Plant pop. pl/ac × 10 ³	SPAD	RLC %	Residue Cover %
	Corn for	Cover	'23 N rate lb/ac	Wet tons ton/ac	Dry tons ton/ac	Moisture %				
1	Grain	None	4.6	12.4 c	4.00 d	67.7 a	33.8	33.1 d	61.7 d	56.0 b
2	Grain	None	180	19.7 a	6.85 abc	65.2 bc	33.9	50.7 ab	94.6 ab	54.5 b
3	Grain	None	220	19.5 ab	6.73 abc	65.5 b	34.2	49.0 b	91.4 b	61.0 ab
4	Silage	None	180	18.7 ab	7.13 ab	61.7 e	33.8	49.9 ab	93.1 ab	19.5 cd
5	Silage	None	220	19.0 ab	6.74 abc	64.6 bcd	34.0	50.1 ab	93.4 ab	26.5 cd
6	Silage	Rye	180	19.6 a	7.29 ab	62.8 de	34.1	49.6 ab	92.5 ab	25.5 cd
7	Silage	Rye	220	19.1 ab	7.30 a	61.9 e	33.9	49.9 ab	93.1 ab	26.5 cd
8	Silage	Blend	180	18.8 ab	6.92 ab	63.3 de	34.2	49.0 b	91.4 b	18.0 d
9	Silage	Blend	220	19.9 a	7.30 a	63.4 cde	34.1	50.6 ab	94.5 ab	28.0 c
10	Grain	None	140	17.8 b	6.15 c	65.4 b	34.0	46.1 c	86.0 c	62.5 ab
11	Grain	None	260	19.4 ab	6.54 bc	66.2 ab	34.1	51.7 a	96.6 a	66.5 a

Stats for RCB Design all 11 treatments

Treatment

P > F:	<0.001	<0.001	<0.001	0.916	<0.001	<0.001	<0.001
LSD 0.10:	1.9	0.74	1.8	NS	2.7	5.0	8.6

Stats for RCB Design with a split-plot arrangement for treatments 2 through 9

Crop and cover crop system

Grain corn, no cover crop	19.6 A	6.79 A	65.4 A	34.1 A	49.8 A	93.0 A	57.8 A
Silage corn, no cover crop	18.8 A	6.93 A	63.1 B	33.9 A	50.0 A	93.3 A	23.0 B
Silage corn, cereal rye	19.4 A	7.29 A	62.3 B	34.0 A	49.7 A	92.8 A	26.0 B
Silage corn, annual blend	19.4 A	7.11 A	63.3 B	34.1 A	49.8 A	92.9 A	23.0 B
P > F:	0.845	0.625	0.033	0.674	0.996	0.997	<0.001

Nitrogen rate for corn in 2022

180 lb/ac	19.2 A	7.05 A	63.2 A	34.0 A	49.8 A	92.9 A	29.4 B
220 lb/ac	19.4 A	7.02 A	63.8 A	34.0 A	49.9 A	93.1 A	35.5 A
P > F:	0.615	0.847	0.278	0.692	0.921	0.876	0.002

Interaction (crop system × N rate)

P > F:	0.563	0.385	0.110	0.566	0.397	0.406	0.291
--------	-------	-------	-------	-------	-------	-------	-------

Table 6b. Corn silage quality parameters, silage nutrient concentrations and nutrient removal in silage as affected by cropping system and nitrogen rates in 2022 (all means are LS means).

Treatments			Silage quality parameters [^]									Nutrient concentration in silage			Nutrient removal in silage		
Trt #	Corn for	Cover	'22 N rate lb/ac	CP % of DM	ADF % of DM	aNDF % of DM	uNDF % of NDF	TTNDFD % of NDF	Starch % of DM	RumenS % of starch	Milk Yield [†] lb/ton	P	K % of DM	S	P ₂ O ₅	K ₂ O	S
1	Grain	None	4.6	4.32 c	26.3 a	44.2 a	12.9 ab	45.1 a	28.6 d	75.6 a	2740 f	0.216 a	0.775 a	0.055 c	36.2 e	68 d	4.0 e
2	Grain	None	180	6.76 a	20.0 cd	35.6 def	10.3 e	44.8 ab	39.3 abc	70.9 bc	3259 abc	0.209 a	0.630 bc	0.068 b	78.1 cd	123 c	11.1 cd
3	Grain	None	220	6.77 a	18.4 d	35.9 def	10.5 de	44.2 ab	38.9 abc	71.1 bc	3250 abc	0.209 a	0.651 bc	0.067 b	81.1 abcd	132 abc	11.3 bc
4	Silage	None	180	7.07 a	20.6 bcd	37.2 cde	10.4 de	43.2 bcd	38.0 abc	69.8 bc	3239 abcd	0.223 a	0.630 bc	0.069 ab	91.9 a	136 abc	12.5 abc
5	Silage	None	220	7.22 a	19.4 cd	33.6 f	9.5 e	43.8 abc	41.3 a	65.8 d	3340 a	0.207 a	0.586 c	0.071 ab	84.4 abcd	125 c	12.6 abc
6	Silage	Rye	180	6.81 a	21.0 bc	38.4 cd	12.2 bc	43.4 abc	36.2 c	71.6 b	3128 cd	0.216 a	0.675 bc	0.069 ab	89.9 ab	148 ab	12.6 abc
7	Silage	Rye	220	6.82 a	22.7 b	39.5 bc	12.4 abc	42.1 cd	35.5 c	70.4 bc	3103 d	0.196 a	0.595 c	0.071 ab	80.4 bcd	128 bc	12.7 ab
8	Silage	Blend	180	7.13 a	21.1 bc	37.4 cde	11.5 cd	43.3 bc	37.9 abc	70.6 bc	3170 bcd	0.221 a	0.634 bc	0.071 ab	91.8 a	138 abc	12.9 ab
9	Silage	Blend	220	7.25 a	19.0 cd	34.6 ef	10.6 de	44.5 ab	40.3 ab	68.4 cd	3287 ab	0.220 a	0.663 bc	0.074 a	91.2 ab	144 abc	13.4 a
10	Grain	None	140	5.66 b	25.2 a	42.7 ab	13.6 a	41.3 d	31.2 d	71.8 b	2923 e	0.194 a	0.663 bc	0.058 c	73.1 d	131 abc	9.5 d
11	Grain	None	260	7.03 a	20.9 bc	37.8 cd	11.8 bc	44.8 ab	36.7 bc	72.4 b	3177 bcd	0.222 a	0.711 ab	0.071 ab	89.4 abc	150 a	12.5 abc

Stats for RCB Design all 11 treatments

Treatment

P > F:	<0.001	<0.001	<0.001	<0.001	0.04	<0.001	0.003	<0.001	0.261	0.068	<0.001	<0.001	<0.001	<0.001
LSD 0.10:	0.50	2.3	3.3	1.2	1.8	3.9	3.1	148	NS	0.090	0.005	11.4	22	1.6

Stats for RCB Design with a split-plot arrangement for treatments 2 through 9

Crop and cover crop system

Grain corn, no cover crop	6.76 A	19.2 B	35.7 A	10.4 BC	44.5 A	39.1 A	71.0 A	3255 A	0.209 A	0.641 A	0.067 A	79.6 B	127 A	11.2 A
Silage corn, no cover crop	7.14 A	20.0 B	35.4 A	10.0 C	43.5 A	39.7 A	67.8 A	3290 A	0.215 A	0.608 A	0.070 A	88.1 A	131 A	12.5 A
Silage corn, cereal rye	6.82 A	21.9 A	39.0 A	12.3 A	42.7 A	35.8 A	71.0 A	3115 B	0.206 A	0.635 A	0.070 A	85.1 AB	138 A	12.7 A
Silage corn, annual blend	7.19 A	20.1 B	36.0 A	11.1 B	43.9 A	39.1 A	69.5 A	3229 A	0.220 A	0.649 A	0.072 A	91.5 A	141 A	13.1 A
P > F:	0.140	0.074	0.118	0.011	0.245	0.234	0.214	0.074	0.303	0.770	0.239	0.089	0.311	0.154

Nitrogen rate for corn in 2022

180 lb/ac	6.94 A	20.7 A	37.2 A	11.1 A	43.7 A	37.8 A	70.7 A	3199 A	0.217 A	0.642 A	0.069 A	87.9 A	136 A	12.3 A
220 lb/ac	7.01 A	19.9 A	35.9 A	10.8 A	43.6 A	39.0 A	68.9 B	3245 A	0.208 A	0.624 A	0.071 A	84.3 A	132 A	12.5 A
P > F:	0.593	0.235	0.194	0.160	0.966	0.288	0.072	0.257	0.121	0.408	0.322	0.234	0.470	0.557

Interaction (crop system × N rate)

P > F:	0.965	0.190	0.245	0.214	0.438	0.455	0.463	0.466	0.501	0.264	0.785	0.426	0.222	0.986
--------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------

[^] CP, crude protein; ADF, acid detergent fiber; aNDF, neutral detergent fiber; uNDF, undigestible NDF at 240 hour; TTNDFD, total tract NDF digestibility; RumenS, in-situ rumen degradable starch at 7 hour.

[†] Milk yield, milk production estimated using the MILK2006 model developed by the Univ. of Wisconsin.

Table 6c. Corn silage quality parameters, silage nutrient concentrations and nutrient removal in silage as affected by cropping system and nitrogen rates in 2023 (all means are LS means).

Trt #	Treatments			Silage quality parameters [^]							Nutrient concentration in silage			Nutrient removal in silage			
	Corn for	Cover	'23 N rate	CP	ADF	aNDF	uNDF	TTNDFD	Starch	RumenS	Milk Yield [†]	P	K	S	P ₂ O ₅	K ₂ O	S
	lb/ac		lb/ac	% of DM	% of DM	% of DM	% of NDF	% of NDF	% of DM	% of starch	lb/ton		% of DM			lb/ac	
1	Grain	None	4.6	4.76 d	30.1 a	51.8 a	15.7 ab	43.1 ab	21.9 d	77.9 a	2516 d	0.222 a	0.848 a	0.055 e	40.7 d	81 d	4.4 f
2	Grain	None	180	6.54 bc	25.1 cde	46.8 cd	14.1 cd	43.0 abc	27.7 abc	74.0 ab	2856 a	0.206 bc	0.744 bcd	0.064 cd	64.6 abc	122 abc	8.7 cd
3	Grain	None	220	6.89 ab	24.7 de	46.3 cd	14.1 cd	43.3 a	28.2 abc	73.6 bc	2855 a	0.212 ab	0.804 ab	0.069 abc	65.5 ab	129 ab	9.3 bcd
4	Silage	None	180	6.90 ab	25.4 cde	46.8 cd	14.7 bc	41.1 de	28.2 abc	69.4 c	2859 a	0.214 ab	0.720 cd	0.069 abc	69.2 a	123 abc	9.7 ab
5	Silage	None	220	7.05 a	25.7 bcde	47.1 cd	14.9 bc	41.4 cde	27.2 bc	69.9 c	2844 a	0.206 bc	0.728 cd	0.071 a	62.9 abc	117 c	9.6 ab
6	Silage	Rye	180	6.23 c	27.9 ab	50.3 ab	16.2 a	39.8 e	24.5 cd	70.7 bc	2704 c	0.197 c	0.710 d	0.062 d	65.7 ab	124 abc	9.0 bcd
7	Silage	Rye	220	6.45 bc	27.1 bc	48.9 abc	15.7 ab	41.0 de	26.4 bc	70.6 bc	2733 bc	0.198 c	0.727 cd	0.065 bcd	66.0 ab	127 abc	9.5 abc
8	Silage	Blend	180	6.63 abc	25.8 bcde	47.9 bc	15.0 bc	41.5 bcd	27.1 bc	70.3 bc	2831 ab	0.209 bc	0.720 d	0.068 abc	65.9 ab	119 bc	9.4 abc
9	Silage	Blend	220	6.79 ab	25.9 bcde	46.8 cd	14.8 bc	41.5 bcd	27.5 abc	70.0 c	2795 abc	0.200 c	0.759 bcd	0.070 ab	66.7 ab	132 a	10.1 a
10	Grain	None	140	6.26 c	24.2 e	44.4 d	13.3 d	42.6 abcd	30.7 a	72.4 bc	2889 a	0.205 bc	0.788 abc	0.062 d	57.9 c	117 c	7.7 e
11	Grain	None	260	6.65 abc	26.5 bcd	48.7 abc	14.6 bc	43.3 a	25.5 bc	73.6 bc	2796 abc	0.204 bc	0.753 bcd	0.065 bcd	60.7 bc	118 c	8.5 d

Stats for RCB Design all 11 treatments

Treatment

P > F:	<0.001	0.008	0.026	0.009	0.016	0.032	0.038	<0.001	0.046	0.026	<0.001	<0.001	<0.001	<0.001
LSD 0.10:	0.46	2.3	3.1	1.2	1.7	3.6	4.0	111	0.012	0.065	0.005	6.9	11	0.8

Stats for RCB Design with a split-plot arrangement for treatments 2 through 9

Crop and cover crop system

Grain corn, no cover crop	6.71 A	24.9 A	46.5 A	14.1 B	43.1 A	28.0 A	73.8 A	2855 A	0.209 A	0.774 A	0.066 A	65.1 A	126 A	9.0 A
Silage corn, no cover crop	6.94 A	25.5 A	46.9 A	14.8 B	41.1 B	27.8 A	69.6 A	2851 A	0.209 A	0.713 A	0.070 A	66.2 A	119 A	9.6 A
Silage corn, cereal rye	6.34 A	27.5 A	49.6 A	15.9 A	40.4 B	25.5 A	70.6 A	2718 B	0.197 A	0.719 A	0.064 A	65.8 A	125 A	9.2 A
Silage corn, annual blend	6.71 A	25.9 A	47.3 A	14.9 AB	41.5 B	27.3 A	70.1 A	2813 AB	0.204 A	0.740 A	0.069 A	66.3 A	126 A	9.8 A
P > F:	0.210	0.166	0.257	0.066	0.054	0.524	0.254	0.096	0.226	0.494	0.253	0.981	0.570	0.343

Nitrogen rate for corn in 2023

180 lb/ac	6.56 B	26.0 A	47.9 A	15.0 A	41.2 A	26.9 A	71.0 A	2812 A	0.206 A	0.718 A	0.065 A	66.4 A	122 B	9.2 B
220 lb/ac	6.79 A	25.8 A	47.3 A	14.8 A	41.8 A	27.3 A	71.0 A	2807 A	0.204 A	0.754 A	0.069 A	65.3 A	126 A	9.6 A
P > F:	0.050	0.712	0.369	0.529	0.240	0.650	0.996	0.820	0.476	0.014	0.025	0.460	0.093	0.039

Interaction (crop system × N rate)

P > F:	0.927	0.822	0.822	0.846	0.772	0.619	0.979	0.827	0.348	0.648	0.778	0.319	0.159	0.435
--------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------

[^] CP, crude protein; ADF, acid detergent fiber; aNDF, neutral detergent fiber; uNDF, undigestible NDF at 240 hour; TTNDFD, total tract NDF digestibility; RumenS, in-situ rumen degradable starch at 7 hour.

[†] Milk yield, milk production estimated using the MILK2006 model developed by the Univ. of Wisconsin.

Table 7a. Monthly mean tile flow, flow-weighted (FW) nitrate-N concentration and nitrate-N loss as affected by treatments in 2022.

Trt #	Corn for	Cover Crop	Total N Rate lb/ac	Tile drain flow								FW nitrate-N concentration								Nitrate-N loss or load							
				April		May		June		Total		April		May		June		Average		April		May		June		Total	
				inch	SE	inch	SE	inch	SE	inch	SE	mg/L	SE	mg/L	SE	mg/L	SE	mg/L	SE	lb/ac	SE	lb/ac	SE	lb/ac	SE	lb/ac	SE
1	Grain	None	4.6	0.8	0.56	2.7	1.15	2.5	0.49	6.2	2.24	3.7	0.6	2.9	1.2	4.1	1.1	3.5	1.1	0.8	0.7	2.8	2.1	2.7	1.1	6.3	3.9
2	Grain	None	180	0.6	0.11	3.3	0.67	2.9	0.61	6.8	1.26	4.4	0.9	4.9	1.3	10.0	1.1	6.3	1.1	0.6	0.1	3.6	0.2	7.1	1.6	11.4	1.6
3	Grain	None	220	1.4	0.51	5.0	1.52	3.3	0.81	9.8	2.77	6.1	0.6	7.2	0.3	12.0	0.6	8.2	0.2	2.1	0.8	9.8	3.6	10.8	3.3	22.7	7.4
4	Silage	None	180	0.7	0.26	3.7	0.33	2.1	0.16	6.6	0.55	6.2	0.2	6.8	0.3	11.3	0.6	8.3	0.2	1.0	0.4	6.4	0.6	5.8	1.0	13.2	0.6
5	Silage	None	220	0.4	0.23	2.7	0.71	1.6	0.24	4.8	1.13	7.3	1.2	9.9	1.6	13.8	1.7	10.5	1.6	0.5	0.2	5.9	0.7	5.8	1.0	12.2	1.5
6	Silage	Rye	180	1.1	0.27	4.2	1.11	2.2	0.48	7.4	1.85	4.5	0.7	3.9	0.5	8.5	0.6	5.3	0.3	1.0	0.4	4.4	1.6	4.8	1.1	10.2	3.0
7	Silage	Rye	220	1.2	0.52	4.6	1.33	3.0	0.47	8.9	2.11	4.3	0.5	5.3	0.3	9.7	1.2	6.1	0.4	1.4	0.6	5.8	1.7	7.6	1.6	14.8	3.1
8	Silage	Blend	180	0.9	0.39	4.4	1.26	2.5	0.56	7.8	2.20	5.2	0.5	5.7	0.8	11.0	1.5	7.4	0.8	1.1	0.4	5.6	1.2	7.0	2.2	13.7	3.7
9	Silage	Blend	220	0.9	0.35	4.4	0.84	2.4	0.35	7.7	1.48	5.5	0.3	6.2	0.5	12.3	0.4	7.9	0.4	1.1	0.4	6.6	1.1	6.9	0.9	14.6	2.3
All treatment mean:				0.9		3.9		2.5		7.3		5.2		5.9		10.3		7.0		1.1		5.7		6.5		13.2	
Fertilized treatment mean:				0.9		4.0		2.5		7.5		5.4		6.2		11.1		7.5		1.1		6.0		7.0		14.1	

Table 7b. Monthly mean tile flow, flow-weighted (FW) nitrate-N concentration and nitrate-N loss as affected by treatments in 2023.

Trt #	Corn for	Cover Crop	Total N Rate lb/ac	Tile drain flow								FW nitrate-N concentration								Nitrate-N loss or load							
				March		April		May		Total †		March		April		May		Average †		March		April		May		Total †	
				inch	SE	inch	SE	inch	SE	inch	SE	mg/L	SE	mg/L	SE	mg/L	SE	mg/L	SE	lb/ac	SE	lb/ac	SE	lb/ac	SE	lb/ac	SE
1	Grain	None	4.6	4.5	2.14	4.1	1.97	5.2	2.15	9.3	4.13	1.4	0.4	1.9	0.4	2.1	0.8	2.1	0.6	1.3	0.5	2.1	1.4	2.9	1.7	5.0	3.0
2	Grain	None	180	5.2	3.55	4.1	0.74	4.7	1.06	8.8	1.79	5.4	1.0	6.7	0.7	6.9	0.9	6.8	0.8	4.2	1.8	6.2	1.2	7.0	1.4	13.1	2.6
3	Grain	None	220	1.1	0.64	5.0	0.66	5.7	0.80	10.8	1.46	9.3	1.1	9.3	1.0	9.6	0.8	9.5	0.7	2.1	1.2	10.9	2.8	12.8	2.6	23.8	5.3
4	Silage	None	180	2.7	1.60	4.0	0.50	4.9	0.78	8.9	1.28	7.3	0.3	8.9	0.7	15.3	2.4	12.5	1.7	4.2	2.2	8.2	1.6	18.1	5.7	26.3	7.3
5	Silage	None	220	5.2	3.94	2.3	1.01	3.8	0.77	6.1	1.56	9.0	0.7	10.0	1.2	15.2	1.3	13.5	0.9	11.8	9.2	5.7	2.6	13.0	2.6	18.7	4.7
6	Silage	Rye	180	1.6	0.77	4.3	1.11	5.4	0.86	9.8	1.96	8.2	0.5	9.5	0.8	13.4	0.8	11.8	0.6	2.8	1.3	9.6	2.6	16.6	3.0	26.2	5.6
7	Silage	Rye	220	1.5	1.02	5.4	1.71	6.3	1.39	11.8	3.12	8.4	0.9	9.2	0.6	14.6	1.4	12.3	1.2	2.4	1.6	11.0	3.7	20.4	4.7	31.5	8.3
8	Silage	Blend	180	0.9	0.49	4.3	0.98	5.7	1.07	10.1	2.04	8.8	0.9	9.0	1.1	12.2	1.2	10.8	1.1	1.5	0.9	8.4	1.8	15.8	3.7	24.2	5.5
9	Silage	Blend	220	0.6	0.53	4.7	0.84	5.9	0.70	10.6	1.52	8.9	0.4	10.8	0.7	15.7	1.4	13.5	0.9	1.2	1.0	11.6	2.5	21.6	4.4	33.2	6.9
All treatment mean:				2.6		4.2		5.3		9.6		7.4		8.4		11.7		10.3		3.5		8.2		14.2		22.5	
Fertilized treatment mean:				2.4		4.3		5.3		9.6		8.1		9.2		12.9		11.3		3.8		9.0		15.7		24.6	

† Annual totals and average does not include tile flow from March 22 through 28, due to erratic flow during that period.

Table 8a. Treatment effects on 3-month flow-weighted nitrate-N concentration, nitrate-N loss and flow-adjusted annual loss in 2022.

Trt #	Corn for	Cover Crop	Total N Rate lb/ac	3-month flow-weighted nitrate-N concentration mg/L				3-month nitrate-N loss lb NO ₃ -N/ac				Flow adj. NO ₃ loss lb/inch	
				Jan-Mar	Apr-Jun	Jul-Sep	Oct-Dec	Average	Jan-Mar	Apr-Jun	Jul-Sep		Oct-Dec
1	Grain	None	4.6	no data	2.7 d	no data	no data	2.7 d	no data	4.3 c	minimal data	4.3 c	0.62 d
2	Grain	None	180		7.9 bc			7.9 bc		11.4 b		11.4 b	1.80 bc
3	Grain	None	220		10.0 ab			10.0 ab		22.7 a		22.7 a	2.25 ab
4	Silage	None	180		9.0 b			9.0 b		13.2 b		13.2 b	2.03 b
5	Silage	None	220		12.5 a			12.5 a		12.2 b		12.3 b	2.82 a
6	Silage	Rye	180		5.9 c			5.9 c		10.2 bc		10.3 bc	1.35 c
7	Silage	Rye	220		7.7 bc			7.7 bc		14.7 b		14.8 b	1.75 bc
8	Silage	Blend	180		8.5 bc			8.5 bc		13.7 b		13.7 b	1.91 bc
9	Silage	Blend	220		8.6 b			8.6 b		14.6 b		14.6 b	1.95 b

Stats for RCB Design all 9 treatments

Treatment

P > F:	0.001	0.001	0.022	0.022	0.001
LSD 0.10:	2.6	2.6	6.6	6.6	0.58

Stats for RCB Design with a split-plot arrangement for treatments 2 through 9

Crop and cover crop system

Grain corn, no cover crop	8.9 AB	8.9 AB	17.0 A	17.0 A	2.02 AB
Silage corn, no cover crop	10.7 A	10.7 A	12.7 A	12.7 A	2.42 A
Silage corn, cereal rye	6.8 C	6.8 C	12.5 A	12.5 A	1.55 C
Silage corn, annual blend	8.5 BC	8.5 BC	14.1 A	14.2 A	1.93 BC
P > F:	0.036	0.035	0.442	0.447	0.038

N rate for corn in 2022

180 lb/ac	7.8 B	7.8 B	12.1 B	12.2 B	1.77 B
220 lb/ac	9.7 A	9.7 A	16.1 A	16.1 A	2.19 A
P > F:	0.026	0.026	0.056	0.056	0.027

Interaction (crop system × N rate)

P > F:	0.485	0.482	0.153	0.154	0.488
--------	-------	-------	-------	-------	-------

Table 8b. Treatment effects on 3-month flow-weighted nitrate-N concentration, nitrate-N loss and flow-adjusted annual loss in 2023.

Trt #	Corn for	Cover Crop	Total N Rate lb/ac	3-month flow-weighted nitrate-N concentration mg/L					3-month nitrate-N loss lb NO ₃ -N/ac					Flow adj. † NO ₃ loss lb/inch
				Jan-Mar	Apr-Jun	Jul-Sep	Oct-Dec	Average†	Jan-Mar	Apr-Jun	Jul-Sep	Oct-Dec	Total †	
1	Grain	None	4.6	1.4 d	2.1 e	no data	no data	2.1 e	1.3	5.0 d	minimal data	5.0 d	0.47 e	
2	Grain	None	180	5.4 c	6.8 d			6.8 d	4.2	13.1 cd		13.1 cd	1.54 d	
3	Grain	None	220	9.6 a	9.5 c			9.5 c	2.1	23.8 ab		23.8 ab	2.15 c	
4	Silage	None	180	7.3 b	12.5 ab			12.5 ab	4.2	26.3 ab		26.3 ab	2.82 ab	
5	Silage	None	220	9.0 ab	13.6 a			13.6 a	11.8	18.7 bc		18.8 bc	3.08 a	
6	Silage	Rye	180	8.2 ab	11.8 ab			11.8 ab	2.8	26.2 ab		26.2 ab	2.67 ab	
7	Silage	Rye	220	8.4 ab	12.4 ab			12.4 ab	2.4	31.5 a		31.6 a	2.80 ab	
8	Silage	Blend	180	8.4 ab	10.9 bc			10.9 bc	1.5	24.2 ab		24.2 ab	2.46 bc	
9	Silage	Blend	220	8.1 ab	13.5 a			13.6 a	1.2	33.2 a		33.3 a	3.07 a	

Stats for RCB Design all 9 treatments

Treatment

P > F:	<0.001	<0.001	<0.001	0.467	0.002	0.002	<0.001
LSD 0.10:	2.0	2.3	2.3	NS	10.2	10.2	0.52

Stats for RCB Design with a split-plot arrangement for treatments 2 through 9

Crop and cover crop system

Grain corn, no cover crop	7.5 A	8.1 B	8.1 B	3.2 A	18.5 A	18.5 A	1.84 B
Silage corn, no cover crop	8.1 A	13.0 A	13.0 A	8.0 A	22.5 A	22.6 A	2.95 A
Silage corn, cereal rye	8.3 A	12.1 A	12.1 A	2.6 A	28.9 A	28.9 A	2.74 A
Silage corn, annual blend	8.2 A	12.2 A	12.2 A	1.3 A	28.7 A	28.7 A	2.76 A
P > F:	0.857	0.003	0.003	0.460	0.120	0.119	0.003

N rate for corn in 2023

180 lb/ac	7.3 B	10.5 B	10.5 B	3.2 A	22.4 A	22.5 A	2.37 B
220 lb/ac	8.8 A	12.2 A	12.2 A	4.4 A	26.8 A	26.9 A	2.77 A
P > F:	0.031	0.026	0.026	0.532	0.191	0.190	0.025

Interaction (crop system × N rate)

P > F:	0.074	0.605	0.608	0.307	0.219	0.221	0.608
--------	-------	-------	-------	-------	-------	-------	-------

† Annual average and total does not include data from March 23 through 28.

Table 10. Soil nitrate-N and ammonium-N from 0- to 30-cm depth as affected by crop system and nitrogen rates for corn.

	Fall 2021				Fall 2022				Fall 2023 [^]			
	4.6†	140	170	LSmean‡	4.6	180	220	LSmean	4.6	180	220	LSmean
----- NO ₃ -N, mg/kg -----												
<u>Cropping system</u>												
Grain, none					2.5	3.2	4.3	3.8 A	1.2	4.2	6.1	5.2 B
Silage, none						9.8	13.5	11.7 A		14.5	25.2	19.8 A
Silage, rye						6.8	15.9	11.4 A		14.5	16.8	15.7 A
Silage, blend						10.6	13.5	12.1 A		16.5	9.5	13.0 AB
LS Mean:						7.6B	11.8A			12.4A	14.4A	
----- NH ₄ -N, mg/kg -----												
<u>Cropping system</u>												
Grain, none					3.4	4.6	3.2	3.9 A	5.3	6.0	6.2	6.1 A
Silage, none						3.9	4.5	4.2 A		5.9	6.8	6.3 A
Silage, rye						4.4	5.0	4.7 A		6.0	6.5	6.2 A
Silage, blend						4.6	4.6	4.6 A		6.6	5.9	6.2 A
LS Mean:						4.4A	4.3A			6.1A	6.3A	

† Control treatment received 4.6 lb N/ac as starter fertilizer.

‡ Numbers followed by different letters are significantly different at $\alpha=0.10$ level. Capital letters signify differences in main effect and small letters are differences due to interaction between main effects.

[^] A significant interaction between main effects was found for NH₄-N in 2023 but is not shown in table (see text).




Figure 1a. Daily precipitation and cumulative tile drainage in 2022.




Figure 1b. Daily precipitation and cumulative tile drainage in 2023.




Figure 2. Inorganic nitrogen in the 0- to 90-cm soil profile on 6 May 2022 as affected by nitrogen rate for corn in 2021.




Figure 3. Inorganic nitrogen in the 0- to 90-cm soil profile on 6 May 2022 as affected by cover crops and N rate for corn in 2021.

Supplemental Materials

Supp. table 1. Experimental methods, products and rates (2021 was a setup year).

Experimental method or procedure	Study year			
	2021	2022	2023	2024
	----- Month / day -----			
Previous crop	Soybean	Corn/silage	Corn/silage	Corn/silage
Total N rates, lb/ac	4.6, 110, 140, 170, 200	4.6, 140, 180, 220, 260	4.6, 140, 180, 220, 260	
Termination herbicides	NA	Roundup PowerMax 24 & 32 oz/ac, Deliver 1 qt/ac	Roundup PowerMax 28 oz/ac, Deliver 1 qt/ac	
Corn hybrid	DeKalb 50-08GENSS RIB	Dairyland HIDF 3802Q	Dairyland HIDF 3802Q	
Pre-emerge herbicides	SureStart 2.5 qt/ac, Roundup WeatherMax 24 oz/ac, Deliver 1 qt/ac	SureStart 2.5 qt/ac	SureStart II 2.5 qt/ac	
Post-emerge herbicides	Liberty 22 oz/ac, Deliver 1 qt/ac	Liberty 32 oz/ac, Class Act Flex (2%)	Liberty 32 oz/ac, Deliver 1 qt/ac	
P-K-S fertilizers sources	TSP ¹ , MOP, Elemental S	TSP, MOP, gypsum	TSP, MOP, gypsum	
P-K-S fertilizer rates, lb/ac	45 lb P ₂ O ₅ , 90 lb K ₂ O/ac, 15 lb S	45 lb P ₂ O ₅ , 90 lb K ₂ O/ac, 15 lb S	45 lb P ₂ O ₅ , 90 lb K ₂ O/ac, 15 lb S	

1 TSP, triple super phosphate (0-46-0); MOP, muriate of potash (0-0-60), elemental S (90%); gypsum (17% S)

Pic. 1. Schematic diagram of tile drainage system.

Pic. 2. Tile drainage well access culvert, data logger, and coolers for holding water sample collection bottles.

Pic. 3. Plumbing inside culvert: sump well, pump, and water meters.

Pic. 4. Silage harvest, cover crop planting (14 September 2021), cover crop growth (November 2021).

Pic. 5. Corn planting on 19 May 2022.

Pic 6. Sidedress UAN application 9 June 2022.

Drainage Research Facility at Waseca, SROC

